



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

[Inclosure.]

Report of infectious disease in Japan, September 23 to September 30, 1897.

Locality.	Cholera.		Dysentery.		Smallpox.	
	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
Kioto Fu.....			63	16		
Osaka Fu.....			51	20		
Tokyo Fu.....	3	2	485	(*)		
Aichi Ken.....			143	43		
Akita Ken.....			11	3		
Awamori Ken.....			166	21		
Chiba Ken.....			177	43		
Fukui Ken.....	(*)					
Fukuoka Ken.....			46	12		
Fukushima Ken.....			102	16	1	
Gifu Ken.....			90	28		
Gumma Ken.....			731	222		
Hiogo Ken.....	(*)					
Hiroshima Ken.....	(*)					
Ibaraki Ken.....			115	37		
Ishikawa Ken.....			46	18		
Iwate Ken.....			30	8		
Kagawa Ken.....			491	145		
Kagoshima Ken.....			86	22		
Kanagawa Ken.....	1	1	310	113		
Kochi Ken.....	(*)					
Kumamoto Ken.....	(*)					
Miyagi Ken.....			106	16	2	
Miyazaki Ken.....			24	5		
Miye Ken.....	(*)					
Nagano Ken.....			1,179	286		
Nagasaki Ken.....	(*)					
Nara Ken.....	(*)					
Niigata Ken.....			1,394	224	5	1
Oita Ken.....	(*)					
Okayama Ken.....			279	70		
Okinawa Ken.....	(*)					
Saga Ken.....			12	4		
Saitama Ken.....			330	166		
Shidzuoka Ken.....			369	118		
Shiga Ken.....			44	9		
Shimane Ken.....			157	52		
Tochigi Ken.....			69	20		
Tokushima Ken.....			140	45		
Tottori Ken.....			204	49		
Toyama Ken.....			12	4		
Wakayama Ken.....			28	7		
Yamagata Ken.....	(*)					
Yamaguchi Ken.....			238	56		
Yamanashi Ken.....			1,251	374		
Yehime Ken.....			137	27		
The Hokkaido.....	(*)					
Taiwan (Formosa).....	(*)					
Total.....	4	3	9,116	2,299	8	1

* No report.

SAN SALVADOR.

*Yellow fever.*SAN SALVADOR, *October 8, 1897.*

SIR: Inclosed find a report made me by Dr. José C. Gasteazoro on the epidemic yellow fever, believing that it may be of interest to the Marine-Hospital Bureau.

May I ask that a copy of the inclosure be sent me, for the reason that a special courier leaves in a few minutes for La Libertad, not giving me the time to make a copy for my archives.

I am, sir, your obedient servant,

JOHN JENKINS,
United States Consul.

Hon. SECRETARY OF THE TREASURY.

[Inclosure.]

*Yellow fever in San Salvador.*SAN SALVADOR, *October 7, 1897.*

The epidemic of yellow fever first started here in the capital in the month of May, 1894. The first person attacked was one of the mail-service boys, who caught it at La Libertad, the port on the Pacific. Ever since we have had the epidemic, but it increases during the months of June, July, August, September, October, and even November; that is, during the rainy season in this country. The cases observed during the months of December, January, February, March, April, and May have been very few and of a very mild character, and generally marked by either influenza or malaria.

We have also noted that the epidemic decreases yearly, both in number and intensity, this applying to both foreigners and natives. This is the general rule of epidemics, and is due to the acclimation of the inhabitants.

In studying yellow fever of Habana, Rio de Janeiro, and New Orleans, and comparing it to ours, we only find differences in the intensity of the symptoms.

Most all our cases that died were on the sixth, seventh, and eighth days, and the causes generally were collapse, intestinal hemorrhage, and uræmia. The mortality during the year of 1894 was 60 per cent; 1895, 40 per cent, and 1896, 38 per cent.

The report for the year 1897 has not been completed. The cases reported now are as follows: July, 38 cases, 11 deaths; August, 54 cases, 16 deaths; September, 34 cases, 12 deaths.

I am, yours, truly,
To Mr. JENKINS,

JOSÉ C. GASTEAZORO, M. D.

United States Consul at San Salvador, Estado Salvador, Rep. Magn. de C. A.

UNITED STATES OF COLOMBIA.

Yellow fever in Cartagena.

[Cablegram.]

CARTAGENA, UNITED STATES OF COLOMBIA, *October 11, 1897.*

Yellow [fever]—SMITH.

Hon. SECRETARY OF STATE.

*Sanitary report from Colon.*COLON, *October 14, 1897.*

SIR: There has been no case of yellow fever at this port for the past month that I am aware of, but in view of the fact that there has been some fever at Bocas del Toro, which is only 140 miles distant, and in constant communication with this port, and the further fact that the fever is very prevalent at Kingston, Jamaica, I suggest that the inspection of vessels clearing for United States ports should be continued for a short while (say a month) longer. * * *

Respectfully, yours,

J. L. PEARCY,

United States Consul.

The SUPERVISING SURGEON-GENERAL,

U. S. Marine-Hospital Service.

STATISTICAL REPORTS.

FRANCE—*Nantes*.—Month of September, 1897. Estimated population, 125,757. Total deaths, 193, including enteric fever, 1; scarlet fever, 1, and diphtheria, 3.

Roubaix.—Month of September, 1897. Estimated population, 130,000. Total deaths, 181, including enteric fever, 2; diphtheria, 2, and whooping cough, 5.